

**Title of Abstract:** Perceived Social Support, Perceived Stress and Stress Symptomatology among Adolescents

**Authors:** Adekeye Olujide Adedapo, Adeusi Sussan Olufunmilola & Akomolafe Adebayo Clement

**Outlet:** *International Journal of Social Sciences and Humanities Review*. Ebonyi State University, Ebonyi, Nigeria, **Vol. 2 (1): 20-29.**

**Date:** 2011

**Abstract:**

This study assessed perceived social support, perceived stress and stress symptomatology among adolescents. A survey was carried out in Covenant University on 237 participants (115 males and 122 females) between ages 14 and 19 years (mean age = 16.8 years). With an aim to verify several relationships, the Multidimensional Scale of Perceived Social Support (Zimet, Dahlem, Zimet & Farley, 1988), Perceived Stress Scale ((PSS), Cohen & Williamson, 1983), and the Stress Symptomatology Scale were administered to the participants. The two hypotheses were sustained. There was a significant combined contribution of perceived social support and perceived stress ( $r = 0.443$ ,  $r^2 = 0.196$ ;  $F_{(2, 234)} = 28.503$ ;  $p < 0.005$ ) on stress symptomatology. There was also a significant difference in stress symptomatology between males and females adolescents ( $t = 3.840$ ;  $df = 235$ ;  $p < 0.05$ ). Recommendations for a re-orientation of adolescents and the use of cognitive-behavioural strategies were made to reduce symptomatic stress reactions among adolescents.

**Keywords:** Perceived stress, Perceived Social support, Stress Symptomatology, adolescents,